

SEQUENTIAL SELECTION OF A MONOTONE SUBSEQUENCE FROM A RANDOM PERMUTATION

PEICHAO PENG AND J. MICHAEL STEELE

(Communicated by David Levin)

ABSTRACT. We find a two term asymptotic expansion for the optimal expected value of a sequentially selected monotone subsequence from a random permutation of length n . A striking feature of this expansion is that it tells us that the expected value of optimal selection from a random permutation is quantifiably larger than optimal sequential selection from an independent sequence of uniformly distributed random variables; specifically, it is larger by at least $(1/6) \log n + O(1)$.

1. SEQUENTIAL SUBSEQUENCE PROBLEMS

In the classical monotone subsequence problem, one chooses a random permutation $\pi : [1 : n] \rightarrow [1 : n]$ and one considers the length of its longest increasing subsequence,

$$L_n = \max\{k : \pi[i_1] < \pi[i_2] < \cdots < \pi[i_k], \text{ where } 1 \leq i_1 < i_2 < \cdots < i_k \leq n\}.$$

On the other hand, in the *sequential* monotone subsequence problem one views the values $\pi[1], \pi[2], \dots$ as though they were presented over time to a decision maker who, when shown the value $\pi[i]$ at time i , must decide (once and for all) either to accept or reject $\pi[i]$ as an element of the selected increasing subsequence.

The decision to accept or reject $\pi[i]$ at time i is based on just the knowledge of the time horizon n and the observed values $\pi[1], \pi[2], \dots, \pi[i]$. Thus, in slightly more formal language, the sequential selection problem amounts to the consideration of random variables of the form

$$(1) \quad L_n^\tau = \max\{k : \pi[\tau_1] < \pi[\tau_2] < \cdots < \pi[\tau_k], \text{ where } 1 \leq \tau_1 < \tau_2 < \cdots < \tau_k \leq n\},$$

where the indices $\tau_i, i = 1, 2, \dots$, are stopping times with respect to the increasing sequence of σ -fields $\mathcal{F}_k = \sigma\{\pi[1], \pi[2], \dots, \pi[k]\}$, $1 \leq k \leq n$. We call a sequence of such stopping times a *feasible selection strategy*, and, if we use τ as a shorthand for such a strategy, then the quantity of central interest here can be written as

$$(2) \quad s(n) = \sup_{\tau} \mathbb{E}[L_n^\tau],$$

where one takes the supremum over all feasible selection strategies.

Received by the editors September 15, 2015 and, in revised form, January 2, 2016 and January 14, 2016.

2010 *Mathematics Subject Classification*. Primary 60C05, 60G40, 90C40; Secondary 60F99, 90C27, 90C39.

Key words and phrases. Monotone subsequence problem, sequential selection, online selection, Markov decision problem, nonlinear recursion, prophet inequality.

It was conjectured in Baer and Brock (1968) that

$$(3) \quad s(n) \sim \sqrt{2n} \quad \text{as } n \rightarrow \infty,$$

and a proof of this relation was first given in Samuels and Steele (1981). A much simpler proof of (3) was later given by Gnedin (2000) who made use of a recursion that had been used for numerical computations by Baer and Brock (1968). The main purpose of this paper is to show how by a more sustained investigation of that recursion one can obtain a two term expansion.

Theorem 1 (Sequential selection from a random permutation). *For $n \rightarrow \infty$ one has the asymptotic relation*

$$(4) \quad s(n) = \sqrt{2n} + \frac{1}{6} \log n + O(1).$$

Given what is known for some closely related problems, the explicit second order term $(\log n)/6$ gives us an unanticipated bonus. For comparison, suppose we consider sequential selection from a sequence of n independently uniformly distributed random variables X_1, X_2, \dots, X_n . In this problem a feasible selection strategy τ is again expressed by an increasing sequence of stopping times $\tau_j, j = 1, 2, \dots$, but now the stopping times are adapted to the increasing σ -fields $\hat{\mathcal{F}}_j = \sigma\{X_1, X_2, \dots, X_j\}$. The analog of (1) is then

$$(5) \quad \hat{L}_n^\tau = \max\{k : X_{\tau_1} < X_{\tau_2} < \dots < X_{\tau_k}, \quad \text{where } 1 \leq \tau_1 < \tau_2 < \dots < \tau_k \leq n\},$$

and the analog of (2) is given by

$$\hat{s}(n) = \sup_{\tau} \mathbb{E}[\hat{L}_n^\tau].$$

Bruss and Robertson (1991) found that for $\hat{s}(n)$ one has a uniform upper bound

$$(6) \quad \hat{s}(n) \leq \sqrt{2n} \quad \text{for all } n \geq 1,$$

so, by comparison with (4), we see there is a sense in which sequential selection of a monotone subsequence from a permutation is *easier* than sequential selection from an independent sequence. In part, this is intuitive; each successive observation from a permutation gives useful information about the subsequent values that can be observed. By (4) one quantifies how much this information helps, and, so far, we have only an analytical understanding of the source of $(1/6) \log n$. A genuinely probabilistic understanding of this term remains elusive.

Since (6) holds for all n and since (4) is only asymptotic, it also seems natural to ask if there is a relation between $\hat{s}(n)$ and $s(n)$ that is valid for all n . There is such a relation if one gives up the logarithmic gap.

Theorem 2 (Selection for random permutations vs. random sequences). *One has for all $n = 1, 2, \dots$ that*

$$\hat{s}(n) \leq s(n).$$

Here we should also note that much more is known about $\hat{s}(n)$ than just (6); in particular, there are several further connections between $s(n)$ and $\hat{s}(n)$. These are taken up in a later section, but first it will be useful to give the proofs of Theorems 1 and 2.

The larger context for the problems studied here is the theory of Markov decision processes (or MDPs) which is closely tied to the theory of optimal stopping and the theory of on-line algorithms (cf. Puterman (1994), Shiryaev (2008), and Flat and Woeginger (1998)). The traditional heart of the theory of MDPs is the optimality equation (or Bellman equation) which presents itself here as the identity (7). One of our main motivations has been the expectation that (7) gives one an appropriate path for examining how one can extract delicate asymptotic information from max-type non-linear recursions of the kind that occur in the theory of MDPs. In this respect, it seems hopeful that tools that parallel the comparison principles of Section 3 and the approximate solutions of Section 4 may be broadly applicable, although the details will necessarily vary from problem to problem.

The proof of Theorem 1 takes most of our effort, and it is given over the next few sections. Section 2 develops the basic recurrence relations, and Section 3 develops stability relations for these recursions. In Section 4 we then do the calculations that support a candidate for the asymptotic approximation of $s(n)$, and we complete the proof of Theorem 1. Our arguments conclude in Section 5 with the brief—and almost computation free—proof of Theorem 2. Finally, in Section 6 we discuss further relations between $s(n)$, $\widehat{s}(n)$, and some other closely related quantities that motivate consideration of two open problems.

2. RECURRENCE RELATIONS

One can get a recurrence relation for $s(n)$ by first step analysis. Specifically, we take a random permutation $\pi : [1 : n + 1] \rightarrow [1 : n + 1]$, and we consider its initial value $\pi[1] = k$. If we reject $\pi[1]$ as an element of our subsequence, we are faced with the problem of sequential selection from the reduced random permutation π' on an n -element set. Alternatively, if we choose $\pi[1] = k$ as an element of our subsequence, we are then faced with the problem of sequential selection for a reduced random permutation π'' of the set $\{k + 1, k + 2, \dots, n + 1\}$ that has $n + 1 - k$ elements. By taking the better of these two possibilities, we get from the uniform distribution of $\pi[1]$ that

$$(7) \quad s(n + 1) = \frac{1}{n + 1} \sum_{k=1}^{n+1} \max\{s(n), 1 + s(n + 1 - k)\}.$$

From the definition (2) of $s(n)$ one has $s(1) = 1$, so subsequent values can then be computed by (7). For illustration and for later discussion, we note that one has the approximate values:

n	1	2	3	4	5	6	7	8	9	10
$s(n)$	1	1.5	2	2.375	2.725	3.046	3.333	3.601	3.857	4.098
$\sqrt{2n}$	1.414	2	2.449	2.828	3.162	3.464	3.742	4	4.243	4.472

Here we observe that for the 10 values in the table one has $s(n) \leq \sqrt{2n}$, and, in fact, this relation persists for all $1 \leq n \leq 174$. Nevertheless, for $n = 175$ one has $\sqrt{2n} < s(n)$, just as (4) requires for all sufficiently large values of n .

We also know from (2) that the map $n \mapsto s(n)$ is strictly monotone increasing, and, as a consequence, the recursion (7) can be written a bit more simply as

$$(8) \quad \begin{aligned} s(n+1) &= \frac{1}{n+1} \max_{1 \leq k \leq n} \left\{ (n-k+1)s(n) + \sum_{i=n-k+1}^n \{s(i) + 1\} \right\} \\ &= \frac{1}{n+1} \max_{1 \leq k \leq n} \left\{ (n-k+1)s(n) + k + \sum_{i=n-k+1}^n s(i) \right\}. \end{aligned}$$

In essence, this recursion goes back to Baer and Brock (1968), p. 408, and it is the basis of most of our analysis.

3. COMPARISON PRINCIPLES

Given a map $g : \mathbb{N} \rightarrow \mathbb{R}$ and $1 \leq k \leq n$, it will be convenient to set

$$(9) \quad H(n, k, g) = k + (n - k + 1)g(n) + \sum_{i=n-k+1}^n g(i),$$

so the optimality recursion (8) can be written more succinctly as

$$(10) \quad s(n+1) = \frac{1}{n+1} \max_{1 \leq k \leq n} H(n, k, s).$$

The next two lemmas make rigorous the idea that if g is almost a solution of (10) for all n , then $g(n)$ is close to $s(n)$ for all n .

Lemma 3 (Upper comparison). *If $\delta : \mathbb{N} \rightarrow \mathbb{R}^+$, $1 \leq g(1) + \delta(1)$, and*

$$(11) \quad \frac{1}{n+1} \max_{1 \leq k \leq n} H(n, k, g) \leq g(n+1) + \delta(n+1) \quad \text{for all } n \geq 1,$$

then one has

$$(12) \quad s(n) \leq g(n) + \sum_{i=1}^n \delta(i) \quad \text{for all } n \geq 1.$$

Proof. We set $\Delta(i) = \delta(1) + \delta(2) + \dots + \delta(i)$, and we argue by induction. Specifically, using (12) for $1 \leq i \leq n$ we have

$$\begin{aligned} H(n, k, s) &= k + (n - k + 1)s(n) + \sum_{i=n-k+1}^n s(i) \\ &\leq k + (n - k + 1)(g(n) + \Delta(n)) + \sum_{i=n-k+1}^n \{g(i) + \Delta(i)\}, \end{aligned}$$

so by monotonicity of $\Delta(\cdot)$ we have

$$\frac{1}{n+1} H(n, k, s) \leq \frac{1}{n+1} H(n, k, g) + \Delta(n).$$

Now, when we take the maximum over $k \in [1 : n]$, the recursion (8) and the induction condition (11) give us

$$\begin{aligned} s(n+1) &\leq \frac{1}{n+1} \max_{1 \leq k \leq n} H(n, k, g) + \Delta(n) \\ &\leq g(n+1) + \delta(n+1) + \Delta(n) = g(n+1) + \Delta(n+1), \end{aligned}$$

so induction establishes (12) for all $n \geq 1$. □

Naturally, there is a lower bound comparison principle that parallels Lemma 3. The statement has several moving parts, so we frame it as a separate lemma even though its proof can be safely omitted.

Lemma 4 (Lower comparison). *If $\delta : \mathbb{N} \rightarrow \mathbb{R}^+$, $g(1) - \delta(1) \leq 1$, and*

$$g(n + 1) - \delta(n + 1) \leq \frac{1}{n + 1} \max_{1 \leq k \leq n} H(n, k, g) \quad \text{for all } n \geq 0,$$

then one has

$$g(n) - \sum_{i=1}^n \delta(i) \leq s(n) \quad \text{for all } n \geq 1.$$

4. AN APPROXIMATION SOLUTION

We now argue that the function $f : \mathbb{N} \rightarrow \mathbb{R}$ defined by

$$(13) \quad f(n) = \sqrt{2n} + \frac{1}{6} \log n$$

gives one an approximate solution of the recurrence equation (8) for $n \mapsto s(n)$.

Proposition 5. *There is a constant $0 < B < \infty$ such that for all $n \geq 1$, one has*

$$(14) \quad -Bn^{-3/2} \leq \frac{1}{n + 1} \left\{ \max_{1 \leq k \leq n} H(n, k, f) \right\} - f(n + 1) \leq Bn^{-3/2}.$$

First step: Localization of the maximum. To deal with the maximum in (14), we first estimate

$$k^*(n) = \text{locmax}_k H(n, k, f).$$

From the definition (9) of $H(n, k, f)$ we find

$$H(n, k + 1, f) - H(n, k, f) = 1 - f(n) + f(n - k),$$

and, from the definition (13) of f , we see this difference is a monotone decreasing function of k ; accordingly, we also have the representation

$$(15) \quad k^*(n) = 1 + \max\{k : 0 \leq 1 - f(n) + f(n - k)\}.$$

Now, for each $n = 1, 2, \dots$ we then consider the function $D_n : [0, n] \rightarrow \mathbb{R}$ defined by setting

$$D_n(x) = 1 - f(n) + f(n - x) = 1 - \{\sqrt{2n} - \sqrt{2(n - x)}\} - \frac{1}{6} \{\log n - \log(n - x)\}.$$

This function is strictly decreasing with $D_n(0) = 1$ and $D_n(n) = -\infty$, so there is a unique solution of the equation $D_n(x) = 0$. For $x \in [0, n)$ we also have the easy bound

$$D_n(x) = 1 - \frac{1}{2} \int_{2(n-x)}^{2n} \frac{1}{\sqrt{u}} du - \frac{1}{6} \log(n/(n - x)) \leq 1 - \frac{x}{\sqrt{2n}}.$$

This gives us $D_n(\sqrt{2n}) \leq 0$, so by monotonicity we have $x_n \leq \sqrt{2n}$.

To refine this bound to an asymptotic estimate, we start with the equation $D_n(x_n) = 0$ and we apply Taylor expansions to get

$$\begin{aligned} 1 &= \sqrt{2n} \left\{ 1 - (1 - x_n/n)^{1/2} \right\} - \frac{1}{6} \log(1 - x_n/n) \\ &= \sqrt{2n} \left\{ \frac{x_n}{2n} + O(x_n^2/n^2) \right\} + O(x_n/n). \end{aligned}$$

By simplification, we then get

$$(16) \quad \sqrt{2n} = x_n + O(x_n^2/n) + O(x_n/n^{1/2}) = x_n + O(1),$$

where in the last step we used our first bound $x_n \leq \sqrt{2n}$.

Finally, by (16) and the characterization (15), we immediately find the estimate that we need for $k^*(n)$.

Lemma 6. *There is a constant $A > 0$ such that for all $n \geq 1$, we have*

$$(17) \quad \sqrt{2n} - A \leq k^*(n) \leq \sqrt{2n} + A.$$

Remark 7. The relations (16) and (17) can be sharpened. Specifically, if we use a two term Taylor series with integral remainders, then one can show $\sqrt{2n} - 2 \leq x_n$. Since we already know that $x_n \leq \sqrt{2n}$, we then see from the characterization (15) and integrality of $k^*(n)$ that we can take $A = 2$ in Lemma 6. This refinement does not lead to a meaningful improvement in Theorem 1, so we omit the details of the expansions with remainders.

Completion of proof of Proposition 5. To prove Proposition 5, we first note that in the definition (9) of $H(n, k, f)$ one has for all $1 \leq k \leq n$ that

$$(18) \quad \frac{1}{n+1}H(n, k, f) = f(n) + \frac{1}{n+1} \left\{ k - \sum_{i=1}^{k-1} (f(n) - f(n-i)) \right\}.$$

The task is to estimate the right-hand side of (18) when $k = k^*(n)$ and $k^*(n)$ is given by (15).

For the moment, we assume that one has $k \leq D\sqrt{n}$, where $D > 0$ is constant. With this assumption, we find that after making Taylor expansions we get from explicit summations that

$$(19) \quad \begin{aligned} \sum_{i=1}^{k-1} (f(n) - f(n-i)) &= \sum_{i=1}^{k-1} \left(\sqrt{2n} - \sqrt{2(n-i)} \right) + \sum_{i=1}^{k-1} \left(\frac{\log n}{6} - \frac{\log(n-i)}{6} \right) \\ &= \sum_{i=1}^{k-1} \left(\frac{i}{\sqrt{2n}} + \frac{i^2}{4n\sqrt{2n}} + O\left(\frac{i^3}{n^{5/2}}\right) \right) + \sum_{i=1}^{k-1} \left(\frac{i}{6n} + O\left(\frac{i^2}{n^2}\right) \right) \\ &= \frac{(k-1)k}{2\sqrt{2n}} + \frac{(k-1)k(2k-1)}{24n\sqrt{2n}} + \frac{(k-1)k}{12n} + O(n^{-1/2}), \end{aligned}$$

where the implied constant of the remainder term depends only on D .

We now define $r(n)$ by the relation $k^*(n) = \sqrt{2n} + r(n)$, and we note by (17) that $|r(n)| \leq A$. Direct algebraic expansions then give us the elementary estimates

$$\frac{(k^*(n) - 1)k^*(n)}{12n} = \frac{1}{6} + O(n^{-1/2})$$

and

$$\frac{(k^*(n) - 1)k^*(n)(2k^*(n) - 1)}{24n\sqrt{2n}} = \frac{1}{6} + O(n^{-1/2}),$$

where in each case the implied constant depends only on A .

Estimation of the first summand of (19) is slightly more delicate than this since we need to account for the dependence of this term on $r(n)$; specifically we have

$$\begin{aligned} \frac{(k^*(n) - 1)k^*(n)}{2\sqrt{2n}} &= \frac{(\sqrt{2n} + r(n) - 1)(\sqrt{2n} + r(n))}{2\sqrt{2n}} \\ &= \sqrt{n/2} + r(n) - \frac{1}{2} + O(n^{-1/2}). \end{aligned}$$

Now, for a pleasing surprise, we note from the last estimate and from the definition of $k^*(n)$ and $r(n)$ that we have cancellation of $r(n)$ when we then compute the critical sum; thus, one has simply

$$(20) \quad k^*(n) - \sum_{i=1}^{k^*(n)-1} (f(n) - f(n - i)) = \sqrt{n/2} + \frac{1}{6} + O(n^{-1/2}).$$

Finally, from the formula (13) for $f(\cdot)$, we have the Taylor expansion

$$(21) \quad f(n + 1) - f(n) = \frac{1}{\sqrt{2n}} + \frac{1}{6n} + O(n^{-3/2}),$$

so, when we return to the identity (18), we see that the estimates (20) and (21) give us the estimate

$$\begin{aligned} &\frac{1}{n + 1} \left\{ \max_{1 \leq k \leq n} H(n, k, f) \right\} - f(n + 1) \\ &= \frac{1}{n + 1} \left(\sqrt{n/2} + \frac{1}{6} + O(n^{-1/2}) \right) + f(n) - f(n + 1) = O(n^{-3/2}). \end{aligned}$$

Here the implied constant is absolute, and the proof of Proposition 5 is complete.

Completion of proof of Theorem 1. Lemmas 3 and 4 combine with Proposition 5 to tell us that by summing the sequence $n^{-3/2}$, $n = 1, 2, \dots$, and by writing $\zeta(z) = 1 + 2^{-z} + 3^{-z} + \dots$ one has

$$|s(n) - f(n)| \leq \zeta(3/2)B \leq (2.62)B \quad \text{for all } n \geq 1.$$

This is slightly more than one needs to complete the proof of Theorem 1.

5. PROOF OF THEOREM 2

The sequential monotone selection problem is a finite horizon Markov decision problem with bounded rewards and finite action space, and for such problems it is known one cannot improve upon an optimal deterministic strategy by the use of strategies that incorporate randomization (cf. Bertsekas and Shreve (1978), Corollary 8.5.1). The proof of Theorem 2 exploits this observation by constructing a randomized algorithm for the sequential selection of a monotone subsequence from a random permutation.

We first recall that if e_i , $i = 1, 2, \dots, n + 1$, are independent exponentially distributed random variables with mean 1 and if one sets

$$Y_i = \frac{e_1 + e_2 + \dots + e_i}{e_1 + e_2 + \dots + e_{n+1}},$$

then the vector (Y_1, Y_2, \dots, Y_n) has the same distribution as the vector of order statistics $(X_{(1)}, X_{(2)}, \dots, X_{(n)})$ of an i.i.d. sample of size n from the uniform distribution (see e.g. Feller (1971), p. 77). Next we let \mathcal{A} denote an optimal algorithm for sequential selection of an increasing subsequence from an independent sample

X_1, X_2, \dots, X_n from the uniform distribution, and we let $\tau(\mathcal{A})$ denote the associated sequence of stopping times. If $\widehat{L}_n^{\tau(\mathcal{A})}$ denotes the length of the subsequence that is chosen from X_1, X_2, \dots, X_n when one follows the strategy $\tau(\mathcal{A})$ determined by \mathcal{A} , then by optimality of \mathcal{A} for selection from X_1, X_2, \dots, X_n we have

$$\widehat{s}(n) = \sup_{\tau} \mathbb{E}[\widehat{L}_n^{\tau}] = \mathbb{E}[\widehat{L}_n^{\tau(\mathcal{A})}].$$

We use the algorithm \mathcal{A} to construct a new randomized algorithm \mathcal{A}' for sequential selection of an increasing subsequence from a random permutation $\pi : [n] \mapsto [n]$. First, the decision maker generates independent exponential random variables e_i , $i = 1, 2, \dots, n + 1$, as above. This is done off-line, and this step can be viewed as an internal randomization.

Now, for $i = 1, 2, \dots, n$, when we are presented with $\pi[i]$ at time i , we compute $X_i = Y_{\pi[i]}$. Finally, if at time i the value X_i would be accepted by the algorithm \mathcal{A} , then the algorithm \mathcal{A}' accepts $\pi[i]$. Otherwise the newly observed value $\pi[i]$ is rejected. By our construction we have

$$(22) \quad \mathbb{E}[L_n^{\tau(\mathcal{A}')}] = \mathbb{E}[\widehat{L}_n^{\tau(\mathcal{A})}] = \widehat{s}(n).$$

Moreover, \mathcal{A}' is a randomized algorithm for the construction of an increasing subsequence of a random permutation π . By definition, $s(n)$ is the expected length of a monotone subsequence selected from a random permutation by an optimal deterministic algorithm, and by our earlier observation, the randomized algorithm \mathcal{A}' cannot do better. Thus, from (22) one has $\widehat{s}(n) \leq s(n)$, and the proof of Theorem 1 is complete.

6. FURTHER DEVELOPMENTS AND CONSIDERATIONS

The uniform upper bound (6) was obtained by Bruss and Robertson (1991) as a consequence of a bound on the expected value of the random variable

$$N(s) = \max \left\{ |A| : \sum_{i \in A} X_i \leq s \right\},$$

where the observations $\{X_i : i \in [1 : n]\}$ have a common continuous distribution with support in $[0, \infty)$. This bound was extended in Steele (2016) to accommodate non-identically distributed random variables, and, as a consequence, one finds some new bounds for the sequential knapsack problems.

On the other hand, this extension does not help one to refine or generalize (6), since, as Coffman et al. (1987) first observed, the sequential knapsack problem and the sequential increasing subsequence problem are equivalent only when the observations are uniformly and identically distributed. Certainly, one may consider the possibility of analogs of (6) for non-identically distributed random variables, but, as even deterministic examples show, the formulation of such analogs is problematical.

Here we should also note that Gnedin (1999) gave a much different proof of (6), and, moreover, he generalized the bound in a way that accommodates random samples with random sizes. More recently, Arlotto et al. (2015) obtained yet another proof of (6) as a corollary to bounds on the *quickest selection problem*, which is an informal dual to the traditional selection problem.

Since the bound of (6) is now well understood from several points of view, it is reasonable to ask about the possibility of some corresponding uniform bound on

$s(n)$. The numerical values that we noted after the recursion (6) and the relation

$$(23) \quad s(n) = \sqrt{2n} + \frac{1}{6} \log n + O(1)$$

from Theorem 1 both tell us that one cannot expect a uniform bound for $s(n)$ that is as simple as that for $\widehat{s}(n)$ given by (6). Nevertheless, numerical evidence suggests that the $O(1)$ term in (23) is always negative. The tools used here cannot confirm this conjecture, but the multiple perspectives available for (6) give one hope.

A closely related issue arises for $\widehat{s}(n)$ when one considers lower bounds. Here the first steps were taken by Bruss and Delbaen (2001) who considered i.i.d. samples of size N_ν where N_ν is an independent random variable with the Poisson distribution with mean ν . If now we write $\widehat{s}(\nu)$ for the expected value of the length of the subsequence selected by an optimal algorithm in the Bruss-Delbaen framework, then they proved that there is a constant $c > 0$ such that

$$\sqrt{2\nu} - c \log \nu \leq \widehat{s}(\nu);$$

moreover, Bruss and Delbaen (2004) subsequently proved that for the optimal feasible strategy $\tau_* = (\tau_1, \tau_2, \dots)$ the random variable

$$\widehat{L}_{N_\nu}^{\tau_*} = \max\{k : X_{\tau_1} < X_{\tau_2} < \dots < X_{\tau_k}, \text{ where } 1 \leq \tau_1 < \tau_2 < \dots < \tau_k \leq N_\nu\}$$

also satisfies a central limit theorem. Arlotto et al. (2015) considered the de-Poissonization of these results, and it was found that one has the corresponding CLT for $\widehat{L}_n^{\tau_*}$ where the sample size n is deterministic. In particular, one has the bounds

$$\sqrt{2n} - c \log n \leq \widehat{s}(n) \leq \sqrt{2n}.$$

Now, by analogy with (23), one strongly expects that there is a constant $c > 0$ such that

$$(24) \quad \widehat{s}(n) = \sqrt{2n} - c \log n + O(1).$$

Still, a proof of this conjecture is reasonably remote, since, for the moment, there is not even a compelling candidate for the value of c .

For a second point of comparison, one can recall the *non-sequential* selection problem where one studies

$$\ell(n) = \mathbb{E}[\max\{k : X_{i_1} < X_{i_2} < \dots < X_{i_k}, 1 \leq i_1 < i_2 < \dots < i_k \leq n\}].$$

Through a long sequence of investigations culminating with Baik et al. (1999), it is now known that one has

$$(25) \quad \ell(n) = 2\sqrt{n} - \alpha n^{1/6} + o(n^{1/6}),$$

where the constant $\alpha = 1.77108\dots$ is determined numerically in terms of solutions of a Painlevé equation of type II. Romik (2014) gives an elegant account of the extensive technology behind (25), and there are interesting analogies between $\ell(n)$ and $s(n)$. Nevertheless, a proof of the conjecture (24) seems much more likely to come from direct methods like those used here to prove (23).

Finally, one should note that the asymptotic formulas for $n \mapsto \ell(n)$, $n \mapsto s(n)$, and $n \mapsto \widehat{s}(n)$ all suggest that these maps are concave, but so far only $n \mapsto \widehat{s}(n)$ has been proved to be concave (cf. Arlotto et al. (2015), p. 3604).

REFERENCES

- Arlotto, A., E. Mossel, and J. M. Steele. 2015. *Quickest online selection of an increasing subsequence of specified size*, to appear in *Random Structures and Algorithms* (eprint arXiv:1412.7985).
- Arlotto, Alessandro, Vinh V. Nguyen, and J. Michael Steele. 2015. *Optimal online selection of a monotone subsequence: a central limit theorem*, *Stochastic Process. Appl.* **125**, no. 9, 3596–3622, DOI 10.1016/j.spa.2015.03.009. MR3357621
- Baer, R. M. and P. Brock. 1968. *Natural sorting over permutation spaces*, *Math. Comp.* **22**, 385–410. MR0228216
- Baik, Jinho, Percy Deift, and Kurt Johansson. 1999. *On the distribution of the length of the longest increasing subsequence of random permutations*, *J. Amer. Math. Soc.* **12**, no. 4, 1119–1178, DOI 10.1090/S0894-0347-99-00307-0. MR1682248
- Bertsekas, Dimitri P. and Steven E. Shreve. 1978. *Stochastic optimal control: The discrete time case*, *Mathematics in Science and Engineering*, vol. 139, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London. MR511544
- Bruss, F. Thomas and Freddy Delbaen. 2001. *Optimal rules for the sequential selection of monotone subsequences of maximum expected length*, *Stochastic Process. Appl.* **96**, no. 2, 313–342, DOI 10.1016/S0304-4149(01)00122-3. MR1865761
- Bruss, F. Thomas and Freddy Delbaen. 2004. *A central limit theorem for the optimal selection process for monotone subsequences of maximum expected length*, *Stochastic Process. Appl.* **114**, no. 2, 287–311, DOI 10.1016/j.spa.2004.09.002. MR2101246
- Bruss, F. Thomas and James B. Robertson. 1991. “Wald’s lemma” for sums of order statistics of i.i.d. random variables, *Adv. in Appl. Probab.* **23**, no. 3, 612–623, DOI 10.2307/1427625. MR1122878
- Coffman, E. G., Jr., L. Flatto, and R. R. Weber. 1987. *Optimal selection of stochastic intervals under a sum constraint*, *Adv. in Appl. Probab.* **19**, no. 2, 454–473, DOI 10.2307/1427427. MR889945
- Feller, William. 1971. *An introduction to probability theory and its applications. Vol. II.*, Second edition, John Wiley & Sons, Inc., New York-London-Sydney. MR0270403
- Flat, A. and G. J. Woeginger. 1998. *Online algorithms. The state of the art*; Papers from the Workshop on the Competitive Analysis of On-line Algorithms held in Schloss Dagstuhl, June 1996; Edited by Amos Fiat and Gerhard J. Woeginger. *Lecture Notes in Computer Science*, vol. 1442, Springer-Verlag, Berlin. MR1673029
- Gnedin, Alexander V. 1999. *Sequential selection of an increasing subsequence from a sample of random size*, *J. Appl. Probab.* **36**, no. 4, 1074–1085. MR1742151
- Gnedin, Alexander V. 2000. *A note on sequential selection from permutations*, *Combin. Probab. Comput.* **9**, no. 1, 13–17, DOI 10.1017/S0963548399004149. MR1751299
- Puterman, Martin L. 1994. *Markov decision processes: discrete stochastic dynamic programming*, *Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics*, John Wiley & Sons, Inc., New York. A Wiley-Interscience Publication. MR1270015
- Romik, D. 2014. *The Surprising Mathematics of Longest Increasing Subsequences*, Cambridge University Press, Cambridge.
- Samuels, Stephen M. and J. Michael Steele. 1981. *Optimal sequential selection of a monotone sequence from a random sample*, *Ann. Probab.* **9**, no. 6, 937–947. MR632967
- Shiryaev, A. N. 2008. *Optimal stopping rules*. Translated from the 1976 Russian second edition by A. B. Aries; Reprint of the 1978 translation. *Stochastic Modelling and Applied Probability*, Vol. 8, Springer-Verlag, Berlin. MR2374974
- Steele, J. M. 2016. *The Bruss-Robertson inequality: Elaborations, extensions, and applications*, to appear in *Mathematica Applicanda (Annales Societatis Mathematicae Polonae Series III)* Volume 44, 2016 (eprint arXiv:1510.00843).

DEPARTMENT OF STATISTICS, THE WHARTON SCHOOL, UNIVERSITY OF PENNSYLVANIA, 3730 WALNUT STREET, PHILADELPHIA, PENNSYLVANIA 19104

E-mail address: ppeichao@wharton.upenn.edu

DEPARTMENT OF STATISTICS, THE WHARTON SCHOOL, UNIVERSITY OF PENNSYLVANIA, 3730 WALNUT STREET, PHILADELPHIA, PENNSYLVANIA 19104

E-mail address: steele@wharton.upenn.edu